

University of Groningen

Polarizations in the Classical Groups

Hesselink, Wim H.

Published in:
Mathematische zeitschrift

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1978

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hesselink, W. H. (1978). Polarizations in the Classical Groups. *Mathematische zeitschrift*, 160(3), 217-234.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Polarizations in the Classical Groups

Wim H. Hesselink

Mathematisch Instituut, Rijksuniversiteit Groningen, Postbus 800,
Groningen, The Netherlands

1. Introduction

1.1. Let G be a connected reductive group over an algebraically closed field k . It has an adjoint action on its Lie algebra \mathfrak{g} . Let P be a parabolic subgroup of G . Let $U(P)$ denote its unipotent radical with Lie algebra $\mathfrak{u}(P)$. An element $x \in \mathfrak{u}(P)$ is called a *Richardson element* of $\mathfrak{u}(P)$ if $\dim(Gx) = 2 \dim(G/P)$.

Lemma. Let $x \in \mathfrak{u}(P)$.

(a) $\dim(Gx) \leq 2 \dim(G/P)$.

(b) x is a Richardson element of $\mathfrak{u}(P)$ if and only if Px is dense in $\mathfrak{u}(P)$ and P contains the identity component $Z_G(x)^0$ of the centralizer $Z_G(x)$.

Proof. This follows from the inequality

$$\begin{aligned} \dim(Gx) &= \dim(G) - \dim(Z_G(x)) \leq \dim(G) - \dim(Z_P(x)) = \\ &= \dim(G/P) + \dim(Px) \leq \dim(G/P) + \dim(\mathfrak{u}(P)) = 2 \dim(G/P). \end{aligned}$$

1.2. Assume that \mathfrak{g} has only finitely many orbits of nilpotent elements. It is known that this assumption is fulfilled if G is a classical group or if the characteristic of k is good enough.

Theorem (a) (Richardson [8, 11] (3.9)). Every parabolic group P in G has Richardson elements in $\mathfrak{u}(P)$.

(b) (Borho-Jantzen [3] (5.18)). Let P_1 and P_2 be parabolic groups in G . For $i = 1, 2$ let $x_i \in \mathfrak{u}(P_i)$ be a Richardson element and let L_i be a Levi factor of P_i . If L_1 is conjugate to L_2 in G then x_1 is conjugate to x_2 under G .

1.3. Conversely, let $x \in \mathfrak{g}$ be nilpotent. A parabolic group P in G is called a *polarization* of x if x is a Richardson element of $\mathfrak{u}(P)$. The set of polarizations of x is denoted by $\text{Pol}(x)$.

Remark. So we only consider polarizations of nilpotent elements. A more general definition leads to questions which are usually first reduced to the nilpotent case, cf. the proof of [3] (5.16). The structure of $\text{Pol}(x)$ is interesting for the infinite-dimensional representation theory, cf. [7] and [1] (6.3).

By (1.1) (b) the “disconnected” centralizer $A(x) := Z_G(x)/Z_G(x)^0$ acts on the set $\text{Pol}(x)$. If $P \in \text{Pol}(x)$ its stabilizer $A_P(x)$ in $A(x)$ is equal to $Z_P(x)/Z_G(x)^0$, since parabolic subgroups are self normalizing, cf. [6] (23.1).

Lemma. *Let \mathcal{P} be a conjugacy class of parabolic subgroups of G . If $\mathcal{P} \cap \text{Pol}(x)$ is non-empty then it is one single orbit under the finite group $A(x)$.*

Proof. Let $P, P' \in \mathcal{P} \cap \text{Pol}(x)$, say $P' = \text{int}(g)P$. Both x and $g^{-1}x$ are Richardson elements of $\mathfrak{u}(P)$. Their dense P -orbits in $\mathfrak{u}(P)$ intersect, so there is $p \in P$ with $px = g^{-1}x$. We have $gp \in Z_G(x)$ and $P' = \text{int}(gp)P$.

Corollary. $\text{Pol}(x)$ is a finite set.

Proof. G has only finitely many conjugacy classes of parabolic subgroups.

1.4. We shall describe $\text{Pol}(x)$ for the classical groups: $G = \text{Gl}(n)$, or $\text{char}(k) \neq 2$ and $G = \text{SO}(n)$ or $G = \text{Sp}(n)$. We verify Theorem (1.2) by hand. So it is natural to describe $\text{Pol}(x)$ in the following steps.

Step 1. Describe the conjugacy classes \mathcal{L} of the Levi factors of the elements of $\text{Pol}(x)$.

Step 2. Given class \mathcal{L} of step 1, determine the classes \mathcal{P} of parabolic subgroups with Levi factor in \mathcal{L} . The number of classes \mathcal{P} associated to \mathcal{L} is denoted by $N_0 = N_0(\mathcal{L})$.

Step 3. Given a class \mathcal{P} of step 2, describe $\mathcal{P} \cap \text{Pol}(x)$ which set is an orbit under $A(x)$, by a representative $P \in \mathcal{P} \cap \text{Pol}(x)$ together with its stabilizer $A_P(x)$ in $A(x)$. The number of elements of $\mathcal{P} \cap \text{Pol}(x)$ is denoted by $N_1 = N_1(\mathcal{P})$. So N_1 is the index of $A_P(x)$ in $A(x)$, or equivalently of $Z_P(x)$ in $Z_G(x)$.

1.5. For the classical groups the finite group $A(x)$ is always abelian. So the stabilizer $A_P(x)$ is independent of the choice of $P \in \mathcal{P} \cap \text{Pol}(x)$. It turns out that $A_P(x)$ only depends on the class \mathcal{L} . More precisely we obtain.

Corollary (7.6). *For $i = 1, 2$ let $P(i) \in \text{Pol}(x)$, say with Levi factor $L(i)$. The following conditions are equivalent:*

- (a) $L(1)$ is conjugate to $L(2)$ in G .
- (b) $A_{P(1)}(x) = A_{P(2)}(x)$ (or equivalently $Z_{P(1)}(x) = Z_{P(2)}(x)$).
- (c) $N_1(P(1)) = N_1(P(2))$.

Note that (b) \Rightarrow (c) is trivial. The implication (a) \Rightarrow (c) holds in general, cf. [4] (7.2). These results lead to the following conjecture for a reductive group G and a nilpotent element $x \in \mathfrak{g}$.

Conjecture. *Let $P(1), P(2) \in \text{Pol}(x)$ have Levi factors $L(1)$ and $L(2)$. Then $L(1)$ is conjugate to $L(2)$ in G if and only if $Z_{P(1)}(x)$ is conjugate to $Z_{P(2)}(x)$ in $Z_G(x)$.*

1.6. The solution of the steps 1 and 2 is due to N. Spaltenstein, cf. [9] Section 5. His methods are sufficient to solve step 3 as well. We give a new proof and more explicit results. Finally we give tables for the groups of type $B_2, B_3, B_4, C_2, C_3, C_4, D_4, D_5, D_6$.

It is a pleasure to express our gratitude to Walter Borho and Hanspeter Kraft for a stimulating correspondence, many discussions and much hospitality.

2. Combinatorial Conventions

\mathbb{N} is the set of natural numbers $1, 2, 3, \dots$. If V is a finite set, its cardinality is denoted by $\#(V)$. If x is a real number $[x]$ denotes the largest integer $\leq x$.

We define a *partition* λ to be a finite subset of \mathbb{N}^2 such that if $(p, q) \in \lambda$ and $i \leq p$ and $j \leq q$ then $(i, j) \in \lambda$. So we identify a partition with the corresponding Young diagram. If λ is a partition we define.

$$|\lambda| = \#(\lambda), \quad \lambda_j = \#\{i | (i, j) \in \lambda\}, \quad \lambda^i = \#\{j | (i, j) \in \lambda\}.$$

A partition λ is determined by each of the non-increasing sequences λ_* and λ^* . Clearly $|\lambda| = \sum_j \lambda_j = \sum_i \lambda^i$. One verifies that

$$\sum_j (j-1) \lambda_j = \sum_i \frac{1}{2} \lambda^i (\lambda^i - 1). \quad (2.1)$$

If (p_1, \dots, p_s) is a finite sequence in \mathbb{N} we define the partition $\lambda = \text{ord}(p_1, \dots, p_s)$ by $\lambda_j = \#\{i | p_i \geq j\}$. Then $\lambda_1 = s$ so that $\lambda^s > 0 = \lambda^{s+1}$. There is a permutation σ of $\{1, \dots, s\}$ such that $\lambda^i = p_{\sigma(i)}$ for $1 \leq i \leq s$. In particular $|\lambda| = \sum p_i$.

For a partition λ we define $N(\lambda)$ to be the number of sequences (p_1, \dots, p_s) with $\lambda = \text{ord}(p_1, \dots, p_s)$. Since the above permutation σ is unique up to multiplication by elements of the stabilizer of $(\lambda^1, \dots, \lambda^s)$ we have

$$N(\lambda) = \lambda_1! \prod_{i \geq 1} ((\lambda_i - \lambda_{i+1})!)^{-1}. \quad (2.2)$$

Let G be a reductive group. We define $b(G) = \dim(G/B)$ where B is some (any) Borel group of G . If P is a parabolic subgroup of G with Levi factor L then we have

$$\dim(G/P) = b(G) - b(L). \quad (2.3)$$

For reference below we note the formulas

$$\begin{aligned} b(\text{Gl}(n)) &= \frac{1}{2}n(n-1) \\ b(\text{Sp}(n)) &= \frac{1}{2}(\frac{1}{2}n(n-1) + [\frac{1}{2}n]) \quad (\text{here } n \text{ is even}) \\ b(\text{SO}(n)) &= \frac{1}{2}(\frac{1}{2}n(n-1) - [\frac{1}{2}n]). \end{aligned} \quad (2.4)$$

3. The General Linear Group $\text{Gl}(n)$

3.1. Let V be a vector space over the algebraically closed field k with $\dim(V) = n$. Let G be $\text{Gl}(V)$. Its Lie algebra \mathfrak{g} is identified with $\text{End}(V)$. Let $x \in \mathfrak{g}$ be nilpotent. The endomorphism x has a Jordan normal form. So there is a partition λ with $|\lambda| = n$ and a *Jordan basis* of V consisting of vectors $e(i, j)$, $(i, j) \in \lambda$, such that $x e(i, j) = e(i-1, j)$ if $i > 1$, and $x e(1, j) = 0$. If $m \geq 0$ then $\dim(\text{Im}(x^m)) = \sum_{i > m} \lambda^i$. So the partition λ of x is unique. G acts on \mathfrak{g} by $\text{Ad}(g)x = g x g^{-1}$. It is clear that nilpotent elements x and y of \mathfrak{g} are conjugate under G if

and only if they have the same partition. The centralizer $Z_G(x)$ of an element $x \in \mathfrak{g}$ is open in the linear space $\{y \in \text{End}(V) | yx = xy\}$, and hence connected.

3.2. A (partial) *flag* F in V is a sequence of subspaces (F_0, \dots, F_s) with $F_0 = 0$, $F_s = V$ and $F_{i-1} \subset F_i$, $F_{i-1} \neq F_i$ for $1 \leq i \leq s$. Its *type* is the sequence (p_1, \dots, p_s) given by $p_i = \dim(F_i/F_{i-1})$. Its *stabilizer* P consists of the $g \in G$ with $gF_i = F_i$ for $0 \leq i \leq s$. Now P is a parabolic subgroup of G . Every parabolic subgroup P of G is the stabilizer of a unique flag. The type of its flag is called the *flag type* of P . Parabolic subgroups of G are conjugate if and only if they have the same flag type.

Let P be the stabilizer of the flag $F = (F_0, \dots, F_s)$. Its unipotent radical consists of the elements $g \in P$ with $(g-1)F_i \subset F_{i-1}$ for $1 \leq i \leq s$. So $\mathfrak{u}(P)$ consists of the $x \in \text{End}(V)$ with $x F_i \subset F_{i-1}$ for $1 \leq i \leq s$. If we choose factors L_i with $F_i = F_{i-1} \oplus L_i$ then

$$L = \prod_{i=1}^s \text{Gl}(L_i) \cong \prod_{i=1}^s \text{Gl}(p_i)$$

is a Levi factor of P . So parabolic subgroups P and Q of G with flag types (p_1, \dots, p_s) and (q_1, \dots, q_t) have conjugate Levi factors if and only if $\text{ord}(p_1, \dots, p_s) = \text{ord}(q_1, \dots, q_t)$, cf. Section 2.

3.3. Theorem. Let $x \in \mathfrak{g}$ be nilpotent with partition λ .

(a) The element x has a polarization P with flag type (p_1, \dots, p_s) if and only if $\lambda = \text{ord}(p_1, \dots, p_s)$. In that case P contains $Z_G(x)$ and is unique.

(b) The element x has $N(\lambda)$ polarizations. All its polarizations have conjugate Levi factors, isomorphic to $\prod_{i \geq 1} \text{Gl}(\lambda^i)$.

Proof. Although this is known, cf. [9] (5.3), we indicate a proof as a preparation for the other classical groups.

Assume $\lambda = \text{ord}(p_1, \dots, p_s)$. We construct a flag $F = (F_0, \dots, F_s)$ with flag type (p_1, \dots, p_s) such that $x F_i \subset F_{i-1}$ for $1 \leq i \leq s$. Put $p = p_1$ and $m = \lambda_p - 1$ and $F_1 = \text{Ker}(x) \cap \text{Im}(x^m)$. If $e(i, j)$, $(i, j) \in \lambda$, is a Jordan basis of V with respect to x , cf.

(3.1), then $F_1 = \sum_{j=1}^p ke(1, j)$. It follows that $\dim(F_1) = p_1$. The induced endomorphism x' of V/F_1 has partition μ given by $\mu_j = \lambda_j - 1$ if $j \leq p$, and $\mu_j = \lambda_j$ if $j > p$. One verifies that $\mu = \text{ord}(p'_1, \dots, p'_{s-1})$ where $p'_i = p_{i+1}$. By induction we may assume the existence of a flag $F' = (F'_0, \dots, F'_{s-1})$ in V/F_1 of type (p'_1, \dots, p'_{s-1}) such that $x' F'_i \subset F'_{i-1}$ for $1 \leq i \leq s-1$. Writing $F'_i = F_{i+1}/F_1$ we obtain a flag F which satisfies the requirements. Let P be the stabilizer of F . Then we have $x \in \mathfrak{u}(P)$. Using the formulas (2.3), (2.4) and (2.1) we obtain

$$2 \dim(G/P) = n^2 - n - 2 \sum (j-1) \lambda_j.$$

It follows that $2 \dim(G/P) = \dim(Gx)$, as calculated for example in [5] (3.8). So P is a polarization. Since $Z_G(x)$ is connected P contains $Z_G(x)$. Alternatively this may be verified by the induction.

By the conjugacy of the parabolic subgroup of a given flag type, it follows that every parabolic subgroup, say with flag type (q_1, \dots, q_t) , is the above constructed polarization of some nilpotent element with partition $\text{ord}(q_1, \dots, q_t)$. By the P -conjugacy of the Richardson elements of $u(P)$ this proves part (a). Part (b) follows immediately. See Section 2 for the definition of $N(\lambda)$.

Remark. In the proof we had to verify (1.2) (a) by hand. The assertion (1.2) (b) follows also.

4. The Symplectic and the (Special) Orthogonal Group

We treat these groups in a unified way, cf. [10] p. 253 and [5] (3.1).

4.1. Conventions. All congruences are modulo 2. We fix a number ε equal to 0 or 1. The field k is algebraically closed and of characteristic $\neq 2$. Let V be a vector space over k with $\dim(V) = n$. Let $\varphi: V \times V \rightarrow k$ be a non-degenerate bilinear form satisfying

$$\varphi(v_1, v_2) = (-1)^\varepsilon \varphi(v_2, v_1) \quad v_1, v_2 \in V.$$

Remark. (a) So $n \equiv 0$ if $\varepsilon = 1$. (b) All forms φ are equivalent.

The group $H = H(V, \varphi)$ consists of the elements $g \in \text{Gl}(V)$ which preserve the form φ . The group $G = G(V, \varphi)$ is the identity component of H . It consists of the elements of H with determinant 1. The common Lie algebra \mathfrak{g} of G and H consists of the elements $x \in \text{End}(V)$ satisfying

$$\varphi(xv_1, v_2) + \varphi(v_1, xv_2) = 0 \quad v_1, v_2 \in V.$$

The groups G and H are reductive of rank $l = [\frac{1}{2}n]$. Their types, Dynkin diagrams and classical names are as follows.

$$\left. \begin{array}{ll} n \not\equiv \varepsilon = 1 & C_l \quad \overset{1}{\circ} - \overset{2}{\circ} - \dots - \overset{l-1}{\circ} \rightleftharpoons \overset{l}{\circ} : \\ n \not\equiv \varepsilon = 0 & B_l \quad \overset{1}{\circ} - \overset{2}{\circ} - \dots - \overset{l-1}{\circ} \rightrightarrows \overset{l}{\circ} \\ n \equiv \varepsilon = 0 & D_l \quad \overset{1}{\circ} - \overset{2}{\circ} - \dots - \overset{l-2}{\circ} \begin{array}{l} \nearrow \overset{l-1}{\circ} \\ \searrow \overset{l}{\circ} \end{array} \end{array} \right\} \begin{array}{l} G = H = \text{Sp}(n) \\ G = \text{SO}(n) \neq H = \text{O}(n). \end{array}$$

4.2. We can choose a basis e_1, \dots, e_n of V such that $\varphi(e_i, e_j) \neq 0$ if and only if $i + j = n + 1$. The torus T in G with weight vectors e_1, \dots, e_n is maximal. Let $\lambda_1, \dots, \lambda_n$ be the corresponding weights. They generate the character group $X(T)$ and are subject to the relations $\lambda_i + \lambda_{n+1-i} = 0$, and $\lambda_{l+1} = 0$ if $n = 2l + 1$.

The root system $\Phi = \Phi(G, T)$ consists of the differences $\lambda_i - \lambda_j$ with $i \neq j$, and $i + j \neq n + 1$ if $\varepsilon = 0$. The simple roots $\alpha_1, \dots, \alpha_l$ are chosen $\alpha_i = \lambda_i - \lambda_{i+1}$ if $1 \leq i < l$, $\alpha_l = \lambda_l - \lambda_{l+1} = (1 + \varepsilon)\lambda_l$ if $n \not\equiv \varepsilon$, and $\alpha_l = \lambda_{l-1} - \lambda_{l+1} = \lambda_{l-1} + \lambda_l$ if $n \equiv \varepsilon = 0$. Their numbers correspond to the numbers in the above Dynkin diagram.

Remark. (a). Assume $n \equiv \varepsilon = 0$. Let $r \in H$ be the reflexion given by $re_i = e_i$ if $i \notin \{l, l+1\}$, and $re_l = e_{l+1}$ and $re_{l+1} = e_l$. This reflexion interchanges the roots α_{l-1}

and α_i . The group H is generated by r and G . (b) If $n \neq \varepsilon = 0$ then H is generated by G and $-\text{id}$, which element usually acts trivial.

4.3. A flag $F = (F_0, \dots, F_s)$ in V is called *isotropic* if $F_i^\perp = F_{s-i}$ for $0 \leq i \leq s$. Let (p_1, \dots, p_s) be its type, cf. (3.2). It satisfies $p_i = p_{s+1-i}$ for $1 \leq i \leq s$. Put $t = [\frac{1}{2}s]$, and $q = 0$ if $s = 2t$, and $q = p_{t+1}$ if $s = 2t + 1$. Clearly $q \equiv n$. One verifies the existence of a basis e_1, \dots, e_n of V such that

$$\varphi(e_i, e_j) \neq 0 \Leftrightarrow i + j = n + 1,$$

$$F_j = \sum k e_i, \quad 1 \leq i \leq \dim(F_j).$$

We assume that this is the basis of (4.2). Consider the full isotropic flag $F' = (F'_0, \dots, F'_n)$ given by $F'_j = \sum_{i=1}^j k e_i$. Let P and B be the stabilizers in G of the flags F and F' , respectively, cf. (3.2). The group B is the Borel group of G corresponding to the basis $\alpha_1, \dots, \alpha_l$ of Φ . So P is some standard parabolic group P_I , cf. [6] (30.1).

We say that the flag F is *not admissible* if there is a P -invariant subspace F' and an index j with $F_j \subset F' \subset F_{j+1}$ and $F_j \neq F' \neq F_{j+1}$.

Lemma. (a) *The flag F is admissible if and only if $\varepsilon = 1$ or $q \neq 2$.*

(b) *Assume that F is admissible. Then $P = P_I$ where I consists of the indices i with $1 \leq i \leq l$ and $i \neq \dim(F_j)$ for all j .*

(c) *Every parabolic subgroup of G is the stabilizer of some admissible isotropic flag.*

Proof. (a) and (b) are left to the reader. (c) Every parabolic subgroup is conjugate to some P_I with $I \subset \{1, \dots, l\}$. By Remark (4.2) (a) we may assume that $n \neq \varepsilon$ or that $l - 1 \in I$ or that $l \notin I$. By (a) all such sets I occur in (b).

4.4. By Lemma (4.3)(b) the type (p_1, \dots, p_s) of an admissible isotropic flag with stabilizer P is a well defined invariant of P . It is called the *flag type* of P .

Lemma. *Parabolic subgroups in G are conjugate under the action of H if and only if they have the same flag type. The class of parabolic subgroups of G with flag type (p_1, \dots, p_s) splits into two conjugacy classes under the action of G if and only if $\varepsilon = 0$ and $s = 2t$ and $p_t \geq 2$.*

4.5. Let $F = (F_0, \dots, F_s)$ be an admissible isotropic flag of type (p_1, \dots, p_s) . Let P be its stabilizer in G . Let L be a Levi factor of P and let S be the radical of L . So S is a maximal torus of the radical of P and L is the centralizer of S in G , cf. [6] (30.2). Consider the weight space decomposition $V = \sum V(\chi)$ of V with respect to S . For $\omega \in X(S)$ we have

$$V(\omega)^\perp = \sum_{\chi \neq -\omega} V(\chi).$$

The group L consists of the $g \in G$ with $gV(\chi) = V(\chi)$ for all $\chi \in X(S)$. One verifies that $X(S)$ has a basis χ_1, \dots, χ_t where $t = [\frac{1}{2}s]$ such that

$$V = V(0) \oplus \sum_{i=1}^t (V(\chi_i) \oplus V(-\chi_i)),$$

$$F_j = \sum_{i=1}^j V(\chi_i) \quad \text{if } j \leq t.$$

We may identify

$$L = G(V(0), \varphi|V(0)) \times \prod_{i=1}^t \text{Gl}(V(\chi_i)).$$

In the notations of (4.3) we have

$$p_i = \dim(V(\chi_i)) \quad \text{if } i \leq t,$$

$$q = \dim(V(0)).$$

We put $v = \text{ord}(p_1, \dots, p_t)$, cf. Section 2. The pair $(v; q)$ is called the *type* of L , or the *Levi type* of P .

Remark. If $\varepsilon = 0$ then $\dim(V(0)) \neq 2$, since $\text{SO}(2)$ is a non-trivial torus, see Lemma (4.3)(a).

4.6. An integer q is called *admissible* if $q \geq 0$, and $q \equiv n$, and $q \neq 2$ if $\varepsilon = 0$. We define $\text{Lev}(n)$ to be the set of pairs $(v; q)$ such that q is admissible and v is a partition with $n = q + 2|v|$.

Lemma. (a) $\text{Lev}(n)$ is the set of the Levi types of the parabolic subgroups of G .

(b) Levi factors of parabolic subgroups of G are conjugate under the action of H if and only if they have the same type.

(c) The class of Levi factors of a given type $(v; q)$ splits into two conjugacy classes under the action of G if and only if $q = \varepsilon = 0$ and $v^i \equiv 0$ for all i .

(d) The number of conjugacy classes \mathcal{P} of parabolic subgroups of G with given Levi type $(v; q)$ is equal to $N(v)$ if $q + \varepsilon \geq 1$, and to $(1 + v_1^{-1}v_2)N(v)$ if $q = \varepsilon = 0$.

Proof. (a) and (b) are immediate. (c) Choose a Levi factor L of type $(v; q)$. The class splits if and only if $H \neq G$ and the normalizer N of L in H is contained in G . The first condition is equivalent to $\varepsilon = 0$. The normalizer N is generated by L and $H(V(0), \varphi|V(0))$ and some standard mappings interchanging weight spaces $V(\pm\chi_i)$, $1 \leq i \leq t$. We have $H(V(0), \varphi|V(0)) \subset G$ if and only if $q = 0$. All interchanging mappings have determinant 1 if and only if $v^i \equiv 0$ for all i .

(d) $N(v)$ is the number of types (p_1, \dots, p_s) of isotropic flags in V such that $v = \text{ord}(p_1, \dots, p_t)$ where $t = [\frac{1}{2}s]$, cf. Section 2. By Lemma (4.4) this settles the case $q + \varepsilon \geq 1$. Assume $q = \varepsilon = 0$. By (4.4) the required number is equal to $2N(v) - N'$ where N' is the number of sequences (p_1, \dots, p_t) with $p_t = 1$ and $v = \text{ord}(p_1, \dots, p_t)$. If $v_1 = v_2$ then $N' = 0$. If $v_1 > v_2$ then $N' = N(\tau)$ where τ is the partition given by $\tau_1 = v_1 - 1$ and $\tau_j = v_j$ for $j \geq 2$. By (2.2) we have

$$N(\tau) = v_1^{-1}(v_1 - v_2)N(v).$$

5. Nilpotent Elements

5.1. Let $x \in \mathfrak{g}$ be nilpotent. By (3.1) there is a unique partition λ and a *Jordan basis* $e(i, j)$, $(i, j) \in \lambda$, such that $xe(i, j) = e(i-1, j)$ if $i > 1$ and $xe(1, j) = 0$. By [10] page 259, this basis can be *normalized* such that we have $\varphi(e(i, j), e(p, q)) \neq 0$ if and only if $i+p = \lambda_j + 1$ and $q = \beta(j)$, where β is some permutation of $\{1, \dots, \lambda^1\}$. By the symmetry relations of (4.1), it follows that $\beta^2 = \text{id}$ and $\lambda_{\beta(j)} = \lambda_j$, and that $\beta(j) \neq j$ whenever $\lambda_j \equiv \varepsilon$. Using Remark (4.1)(b) one verifies that any permutation β satisfying these conditions can be used for x .

Let $\text{Pan}(n, \varepsilon)$ denote the set of the partitions λ such that $|\lambda| = n$ and that the number $\#\{j | \lambda_j = m\}$ is even for every $m \in \mathbb{N}$ with $m \equiv \varepsilon$. Again using Remark (4.1)(b) we obtain

Lemma. (a) *A partition λ is the partition of some nilpotent element $x \in \mathfrak{g}$ if and only if $\lambda \in \text{Pan}(n, \varepsilon)$.*

(b) *Nilpotent elements x and y of \mathfrak{g} are conjugate under H if and only if they have the same partition.*

5.2. Let $x \in \mathfrak{g}$ be nilpotent with partition λ and normalized Jordan basis as in (5.1). We put $M_i = \text{Ker}(x) \cap \text{Im}(x^{i-1})$. The vectors $e(1, j)$ with $1 \leq j \leq \lambda^i$ form a basis of M_i , so that $\dim(M_i) = \lambda^i$. One verifies the existence of a unique bilinear form φ_i on M_i such that $\varphi_i(v, v') = \varphi(v, v'')$ if $v, v' \in M_i$ and $v' = x^{i-1}v''$. This form satisfies

$$\varphi_i(v_1, v_2) = (-1)^{i-1+\varepsilon} \varphi_i(v_2, v_1) \quad \text{if } v_1, v_2 \in M_i.$$

One verifies that M_{i+1} is the kernel of the form φ_i on M_i . So there is an induced non-degenerate bilinear form φ'_i on M_i/M_{i+1} . Let $C_i = H(M_i/M_{i+1}, \varphi'_i)$ be the corresponding orthogonal or symplectic group, cf. (4.1). The canonical morphism

$$Z_H(x) \rightarrow \prod_i C_i$$

is surjective. Its kernel is connected. If $\lambda^i = \lambda^{i+1}$ or $i \equiv \varepsilon$, the group C_i is connected. If $\lambda^i > \lambda^{i+1}$ and $i \not\equiv \varepsilon$ then C_i is an orthogonal group with two connected components. Compare [10] page 261.

5.3. Consider the finite groups $A(x) = Z_G(x)/Z_G(x)^0$ and $A'(x) = Z_H(x)/Z_H(x)^0$, cf. (1.3). We have $A(x) \subset A'(x)$. By (5.2) the group $A'(x)$ is abelian and all its elements have order 2. We may consider $A'(x)$ as a vector space over \mathbb{F}_2 with a basis indexed by the pairs (i, λ^i) such that $\lambda^i > \lambda^{i+1}$ and $i \not\equiv \varepsilon$. This index set is equal to the set of pairs (λ_j, j) such that

$$j \in B(\lambda) = \{j \in \mathbb{N} | \lambda_j > \lambda_{j+1}, \lambda_j \not\equiv \varepsilon\}.$$

So we may identify

$$A'(x) = \{a \in \mathbb{F}_2^{\mathbb{N}} | j \notin B(\lambda) \Rightarrow a_j = 0\}.$$

If $\varepsilon=1$ then $G=H$ so that $A(x)=A'(x)$. If $\varepsilon=0$ the group G consists of the elements $g \in H$ with determinant 1. In general we obtain

$$A(x) = \{a \in \mathbb{F}_2^{\mathbb{N}} \mid j \notin B(\lambda) \Rightarrow a_j = 0; \varepsilon = 0 \Rightarrow \sum a_j = 0\}.$$

Remark. Assume, as we may, that the normalization of (5.1) satisfies $\beta(j)=j$ if and only if $\lambda_j \neq \varepsilon$. For $m \in \mathbb{N}$ with $\lambda_m \neq \varepsilon$ we define the reflexion $r(m) \in Z_H(x)$ by

$$\begin{aligned} r(m)e(i, j) &= e(i, j) & \text{if } m \neq j, \\ r(m)e(i, j) &= -e(i, j) & \text{if } m = j. \end{aligned}$$

The image of $r(m)$ in $A'(x)$ is the vector $r'(m)$ with $r'(m)_j = 1$ if and only if $\lambda_m = \lambda_j > \lambda_{j+1}$. So the vectors $r'(j)$ with $j \in B(\lambda)$ form a basis of $A'(x)$. Compare [9] (3.2), (3.9), (3.14).

5.4. Corollary. *Let $x \in \mathfrak{g}$ be nilpotent with partition λ . The G -orbit of x is unequal to the H -orbit of x if and only if $\varepsilon=0$ and $B(\lambda)$ is empty. In that case the H -orbit of x consists of two G -orbits.*

Proof. This is the only case where $G \neq H$ and $A(x)=A'(x)$. Compare [10] page 264.

6. Combinatorics

6.1. Let $\text{Pai}(n, q)$ denote the set of partitions π such that $\pi_j \equiv 1$ if $j \leq q$, and $\pi_j \equiv 0$ if $j > q$. We define the set $\text{Pai}(n)$ to be the (clearly disjoint) union of the sets $\text{Pai}(n, q)$ with q admissible, cf. (4.6). We identify the set $\text{Lev}(n)$ of (4.6) with $\text{Pai}(n)$ using the bijection $(v; q) \mapsto \pi$ given by

$$\pi_j = 2v_j + 1 \quad \text{if } j \leq q, \quad \pi_j = 2v_j \quad \text{if } j > q. \quad (*)$$

So we write $(v; q) = \pi$ if $(*)$ holds.

Remark. If P is a parabolic group with flag type (p_1, \dots, p_s) , cf. (4.4), we have thus identified its Levi type $(v; q)$ with the partition $\pi = \text{ord}(p_1, \dots, p_s)$.

6.2. The set $\text{Pan}(n, \varepsilon)$ is defined in (5.1). We define the Spaltenstein mapping $S: \text{Pai}(n) \rightarrow \text{Pan}(n, \varepsilon)$ as follows. If $\pi \in \text{Pai}(n)$, put

$$I(\pi) = \{j \in \mathbb{N} \mid j \neq n, \pi_j \equiv \varepsilon, \pi_j \geq \pi_{j+1} + 2\}$$

and let the partition $\lambda = S(\pi)$ be given by

$$\begin{aligned} \lambda_j &= \pi_j - 1 & \text{if } j \in I(\pi), \\ \lambda_j &= \pi_j + 1 & \text{if } j-1 \in I(\pi), \\ \lambda_j &= \pi_j & \text{otherwise.} \end{aligned}$$

It is clear that λ is a partition with $|\lambda| = n$. In (6.4) we verify that $\lambda \in \text{Pan}(n, \varepsilon)$.

6.3. Lemma. *Let $\pi \in \text{Pai}(n, q)$, q admissible, and $\lambda = S(\pi)$.*

(a) *Assume $\lambda_j \neq \pi_j$. Then $\lambda_j = \pi_j + (-1)^{n-j}$. We have $j \leq q$ if $\varepsilon = 1$, and $j > q$ if $\varepsilon = 0$. In either case $\pi_j \equiv \varepsilon \neq \lambda_j$.*

(b) *$\# \{j | \lambda_j \equiv 1\} = q + 2(-1)^{\varepsilon} \# I(\pi)$.*

(c) *If $j \neq n$ and $\lambda_j > \lambda_{j+1}$ then $\lambda_j \equiv \lambda_{j+1} \neq \varepsilon$; if $n \equiv 1$ then $\lambda_1 \neq \varepsilon$.*

Proof. (a) Since $q \equiv n$ we have $q \notin I(\pi)$. (b) follows from (a).

(c) Let $j \neq n$ and $\lambda_j > \lambda_{j+1}$. It follows from (a) that $\pi_j > \pi_{j+1}$. Since $j \neq q$ we have $\pi_j \equiv \pi_{j+1}$, so that $\pi_j \geq \pi_{j+1} + 2$. If $\pi_j \equiv \varepsilon$ then $j \in I(\pi)$ and the assertion follows. If $\pi_j \neq \varepsilon$ then $j \notin I(\pi)$ and again the assertion follows. Assume $n \equiv 1$. Then $\varepsilon = 0$ by Remark (4.1)(a), and $q \neq 0$. It follows that $\pi_1 \equiv 1$ and that $\lambda_1 = \pi_1$.

6.4. For a partition λ with $|\lambda| = n$ we consider the condition $C(\lambda)$: if $j \neq n$ then $\lambda_j \equiv \lambda_{j+1}$.

Lemma. *The assertion (6.3)(c) holds if and only if $C(\lambda)$ holds and $\lambda \in \text{Pan}(n, \varepsilon)$.*

The proof may be left to the reader.

6.5. For a partition λ with $|\lambda| = n$ we define

$$J(\lambda) = \{j | \lambda_j \equiv \varepsilon\} \cup \{j, j+1 | j \equiv n, \lambda_j = \lambda_{j+1}\},$$

$$j_1(\lambda) = \sup \{j \in J(\lambda) | \lambda_j \equiv 1\} \quad (\text{possibly } -\infty),$$

$$j_0(\lambda) = \min \{j \in J(\lambda) | \lambda_j \equiv 0\}.$$

Proposition. *Let q be an admissible integer and $\lambda \in \text{Pan}(n, \varepsilon)$.*

(a) *The restriction $S: \text{Pai}(n, q) \rightarrow \text{Pan}(n, \varepsilon)$ is injective.*

(b) *We have $\lambda \in S(\text{Pai}(n, q))$ if and only if $C(\lambda)$ holds and $j_1(\lambda) \leq q < j_0(\lambda)$.*

(c) *We have $\lambda \in S(\text{Pai}(n))$ if and only if $C(\lambda)$ holds and $j_1(\lambda) < j_0(\lambda)$.*

Proof. (a) follows from (6.3)(a).

(b) Assume $\lambda = S(\pi)$ with $\pi \in \text{Pai}(n, q)$. Condition $C(\lambda)$ holds by (6.3)(c) and (6.4). If $j \equiv n$ and $\lambda_j = \lambda_{j+1}$, then $\lambda_j \geq \pi_j \geq \pi_{j+1} \geq \lambda_{j+1}$ so that $\lambda_j = \pi_j$ and $\lambda_{j+1} = \pi_{j+1}$. Together with (6.3)(a) this proves that $\lambda_j = \pi_j$ whenever $j \in J(\lambda)$. This implies $j_1(\lambda) \leq q < j_0(\lambda)$.

Assume that $C(\lambda)$ holds and that $j_1(\lambda) \leq q < j_0(\lambda)$. Let π_* be the unique sequence in \mathbb{Z} with $\pi_j = \lambda_j - (-1)^{n-j} \delta_j$ where $\delta_j \in \{0, 1\}$ such that $\pi_j \equiv 1$ if and only if $j \leq q$. The inequality implies that $\pi_j = \lambda_j$ whenever $j \in J(\lambda)$.

If $\lambda_j \geq \lambda_{j+1} + 2$ then $\pi_j \geq \pi_{j+1}$. If $\lambda_j = \lambda_{j+1} + 1$, then $j \in J(\lambda)$ or $j+1 \in J(\lambda)$, so that $\pi_j \geq \pi_{j+1}$. If $j \equiv n$ and $\lambda_j = \lambda_{j+1}$, then $j \in J(\lambda)$ and $j+1 \in J(\lambda)$, so that $\pi_j = \pi_{j+1}$. If $j \neq n$ then $\pi_j \geq \lambda_j \geq \lambda_{j+1} \geq \pi_{j+1}$. So the sequence π_* is non-increasing. If $\lambda_j = 0$ then $\pi_i = 0$ for all $i > j$. This proves that π_* defines a partition π .

If $j \neq n$ then $\lambda_j \equiv \lambda_{j+1}$ and $\pi_j \equiv \pi_{j+1}$, by $C(\lambda)$ and $q \equiv n$. It follows that $\pi_j = \lambda_j + 1$ if and only if $\pi_{j+1} = \lambda_{j+1} - 1$. Using that $n \equiv 1$ implies $\varepsilon = 0$, one proves that $\pi_1 = \lambda_1 - 1$. It follows that $|\pi| = |\lambda| = n$, so that $\pi \in \text{Pai}(n, q)$.

Moreover it now suffices to prove that $j \in I(\pi)$ if and only if $\pi_j = \lambda_j + 1$. By the last paragraph, $\pi_j = \lambda_j + 1$ implies $j \in I(\pi)$. Assume $j \in I(\pi)$ and $\pi_j \neq \lambda_j + 1$. We have $j \neq n$ and $\pi_j \equiv \varepsilon$ and $\pi_j \geq \pi_{j+1} + 2$ and $\pi_j = \lambda_j$ and $\pi_{j+1} = \lambda_{j+1}$. Condition (6.3)(c) holds by (6.4), so that $\pi_j = \lambda_j \neq \varepsilon$, a contradiction.

(c) Assume that $C(\lambda)$ holds and that $j_1(\lambda) < j_0(\lambda)$. It remains to show the existence of an admissible integer q with $j_1(\lambda) \leq q < j_0(\lambda)$. If $\varepsilon = 1$ then $n \equiv 0$ and either $j_1(\lambda) = -\infty$ or $j_1(\lambda) \equiv n$ by (6.3)(c), so that we may use $q = 0$ or $q = j_1(\lambda)$ respectively. Assume $\varepsilon = 0$. By (6.3)(c) we have $j_0(\lambda) \not\equiv n$. If $j_0(\lambda) \neq 3$ we may use $q = j_0(\lambda) - 1$. If $j_0(\lambda) = 3$ then $n \equiv 0$ and $j_1(\lambda) = -\infty$ so that we may use $q = 0$.

6.6. Lemma. Assume $\varepsilon = 0$. Let $(v; q) = \pi \in \text{Pai}(n)$ and $\lambda = S(\pi)$.

(a) The set $B(\lambda)$ of (5.3) is empty if and only if $q = 0$ and $\lambda = \pi$, if and only if $q = 0$ and $v^i \equiv 0$ for all i .

(b) We have $q > 0$ if and only if there is $j \in B(\lambda)$ such that $j \notin I(\pi)$ and $j - 1 \notin I(\pi)$.

The proof is left to the reader.

7. The Main Theorem

7.1. Let $x \in \mathfrak{g}$ be nilpotent with partition λ , cf. (5.1). We use the description of its disconnected centralizer $A(x)$ given in (5.3). Let $\pi = (v; q)$ be an element of $\text{Pai}(n)$, cf. (6.1). Let $\text{Pol}(x, \pi)$ denote the set of polarizations P of x with Levi type $(v; q)$. The Spaltenstein mapping S is defined in (6.2) and analysed in (6.5). If $\lambda = S(\pi)$, we put

$$u = \# I(\pi) = \frac{1}{2}(-1)^\varepsilon(-q + \# \{j | \lambda_j \equiv 1\}), \quad \text{cf. (6.3)(b).}$$

Theorem. (a) $\text{Pol}(x, \pi)$ is non-empty if and only if $\lambda = S(\pi)$.

(b) If $P \in \text{Pol}(x, \pi)$ its stabilizer $A_P(x)$ in $A(x)$ consists of the elements $a \in A(x)$ with $a_j = a_{j+1}$ whenever $j \in I(\pi)$.

(c) Levi factors of elements of $\text{Pol}(x, \pi)$ are conjugate in G .

(d) If $\lambda = S(\pi)$ then $\text{Pol}(x, \pi)$ consists of N_0 conjugacy classes under $A(x)$ each consisting of N_1 polarizations where

$$N_0 = N(v) \quad \text{and} \quad N_1 = 2^u \quad \text{if} \quad q + \varepsilon \geq 1 \quad \text{or} \quad B(\lambda) \text{ is empty.}$$

$$N_0 = (1 + v_1^{-1} v_2) N(v) \quad \text{and} \quad N_1 = 2^{u-1} \quad \text{if} \quad q = \varepsilon = 0 \quad \text{and} \quad B(\lambda) \neq \emptyset.$$

7.2. Lemma. Let P be a parabolic subgroup with Levi type $(v; q) = \pi$. Assume that $\lambda = S(\pi)$. Then $\dim(Gx) = 2 \dim(G/P)$.

Proof. Let L be a Levi factor of P . By (2.3), (2.4) and (4.6) we have $2 \dim(G/P) = 2b(G) - 2b(L)$ where

$$\begin{aligned} 2b(L) &= \sum v^i(v^i - 1) + \frac{1}{2}q(q-1) - (-1)^\varepsilon \left[\frac{1}{2}q \right] \\ &= \sum \frac{1}{2} \pi^i(\pi^i - 1) - (-1)^\varepsilon \left[\frac{1}{2}q \right]. \end{aligned}$$

By [5] (3.8) we have $\dim(Gx) = 2b(G) - \gamma_\varepsilon(\lambda)$ where

$$\gamma_\varepsilon(\lambda) = \sum (j-1)\lambda_j - (-1)^\varepsilon \left[\frac{1}{2} \# \{j | \lambda_j \equiv 1\} \right].$$

Now the result follows from (2.1) and (6.3)(b).

7.3. Lemma. Assume that $\lambda = S(\pi)$. Let (p_1, \dots, p_s) be a sequence in \mathbb{N} with $\pi = \text{ord}(p_1, \dots, p_s)$ and $p_i = p_{s+1-i}$ for $1 \leq i \leq s$. Then x has a polarization P with flag type (p_1, \dots, p_s) .

Proof. It suffices to construct an isotropic flag $F = (F_0, \dots, F_s)$ of type (p_1, \dots, p_s) such that $x F_i \subset F_{i-1}$ for $1 \leq i \leq s$. In fact by the assumption $\pi \in \text{Pai}(n)$ the flag F will be admissible. So its stabilizer P in G is parabolic with flag type (p_1, \dots, p_s) . The condition $x F_i \subset F_{i-1}$ implies that $x \in \mathfrak{u}(P)$, so P is a polarization of x by (7.2).

Assume $s=1$. Then $\pi_j \leq 1$ for all j . So $I(\pi)$ is empty and $\lambda = \pi$ and $x=0$. We choose $F_0=0$, $F_1=V$.

Assume $s>1$. Put $p=p_1=p_s$. The type (p_2, \dots, p_{s-1}) of the required isotropic flag $(F_i/F_1)_{0 < i < s}$ in F_{s-1}/F_1 gives rise to a partition $\rho \in \text{Pai}(n-2p, q)$ given by

$$\rho_j = \# \{i | 2 \leq i \leq s-1, p_i \geq j\},$$

so that $\rho_j = \pi_j - 2$ if $j \leq p$, and $\rho_j = \pi_j$ if $j > p$.

Let $S: \text{Pai}(n-2p) \rightarrow \text{Pan}(n-2p, \varepsilon)$ be the Spaltenstein mapping, cf. (6.2). Put $\mu = S(\rho)$. There are two cases.

(A) $I(\pi) = I(\rho)$. Then $p \equiv n$ or $\pi_p \not\equiv \varepsilon$ or $\pi_p \geq \pi_{p+1} + 4$. We have $\mu_j = \lambda_j - 2$ if $j \leq p$, and $\mu_j = \lambda_j$ if $j > p$. Note that $\lambda_p > \lambda_{p+1}$ since μ is a partition.

(B) $I(\pi) = \{p\} \cup I(\rho)$ and $p \notin I(\rho)$. Then $p \not\equiv n$, $\pi_p \equiv \varepsilon$ and $\pi_p = \pi_{p+1} + 2$. We have $\mu_j = \lambda_j - 2$ if $j < p$, and $\mu_j = \lambda_j$ if $j > p+1$, and $\mu_p = \lambda_p - 1$, $\mu_{p+1} = \lambda_{p+1-1}$. Moreover $\lambda_p = \lambda_{p+1} \not\equiv \varepsilon$ and $\lambda_{p+1} > \lambda_{p+2}$.

We choose a normalized Jordan basis $e(i, j)$, $(i, j) \in \lambda$, of V with respect to x , cf. (5.1). In case (B) we require moreover that $\beta(p) = p+1$. We put $F_0=0$, $F_s=V$. In case (A) we define $F_1 = \sum_{j=1}^p k e(1, j)$. In case (B) we have two possibilities

$$F_1 = \sum_{j=1}^p k e(1, j) \quad \text{or} \quad F_1 = \sum_{j=1, j \neq p}^{p+1} k e(1, j)$$

In both cases we put $F_{s-1} = F_1^\perp$.

One verifies that $x F_1 = 0$ and hence $x F_s \subset F_{s-1}$, that F_1 is contained in F_{s-1} , and that the induced endomorphism x_1 of F_{s-1}/F_1 has partition μ . We proceed by induction.

7.4. Lemma. Let P be the polarization of x constructed in (7.3). The stabilizer $A_P(x)$ consists of the elements $a \in A(x)$ with $a_j = a_{j+1}$ whenever $j \in I(\pi)$.

Proof. Let F be the flag constructed in (7.3). The group $Z_H(x)^0$ is contained in P and therefore it leaves F invariant. So we may define the stabilizer $A'_F(x)$ of F in $A'(x)$. We have $A_P(x) = A(x) \cap A'_F(x)$.

It suffices to prove that $a \in A'(x)$ leaves F invariant if and only if $a_j = a_{j+1}$ whenever $j \in I(\pi)$. The case $s=1$ is trivial since $I(\pi)$ is empty and $A'(x) = A'_F(x)$. We proceed by induction as in (7.3). Clearly any element of $A'(x)$ leaves F_0 and F_s invariant.

Assume $s>1$. Put $p=p_1=p_s$ and $h=\lambda_p$. In case (A) we have $F_1 = M_h$, cf. (5.2). In case (B) the space F_1 is one of the two hyperplanes in M_h with $\varphi_h|_{F_1} = 0$. It follows that $a \in A'(x)$ leaves F_1 and F_{s-1} invariant if and only if case (A) applies or $a_{p+1} = 0$ and case (B) applies.

Assume that $a \in A'(x)$ leaves F_1 and F_{s-1} invariant. Then a has an image $a' \in A'(x_1)$ where x_1 is the induced endomorphism of F_{s-1}/F_1 . The element a leaves F invariant if and only if a' leaves the induced flag $(F_i/F_1)_{0 < i < s}$ in F_{s-1}/F_1 invariant. By induction this is the case if and only if $a'_j = a'_{j+1}$ whenever $j \in I(\rho)$.

Case (A₁). $I(\pi) = I(\rho)$ and $B(\lambda) = B(\mu)$, so that $\lambda_p \equiv \varepsilon$ or $\lambda_p > \lambda_{p+1} + 2$. After the identifications of (5.3) the map $A'(x) \rightarrow A'(x_1)$ is the identity. One verifies this using the basis of Remark (5.3). Now the assertion follows by induction.

Case (A₂). $I(\pi) = I(\rho)$ and $B(\lambda) = \{p\} \cup B(\mu)$ where $p \notin B(\mu)$. So $\lambda_p \not\equiv \varepsilon$ and $\lambda_p = \lambda_{p+1} + 2$. First assume that $\lambda_p > 2$. We have a unique $i > p$ such that $\mu_p = \mu_{p+1} = \mu_i > \mu_{i+1}$. Moreover $i \in B(\mu)$. In the notation of Remark (5.3) we have that the images of $r(p)$ and $r(i)$ in $A'(x_1)$ are both equal to $r'(i)$. It follows that $a' \in A'(x_1)$ is given by $a'_j = a_j$ if $j \notin \{p, i\}$ and $a'_p = 0$ and $a'_i = a_p + a_i$. Now it suffices to verify that $a_j - a_{j+1} = a'_j - a'_{j+1}$ whenever $j \in I(\rho)$. In fact $I(\pi) = I(\rho)$, so we may proceed by induction. Let $j \in I(\rho)$. Then we have $\mu_j \geq \mu_{j+1} > \mu_{j+2}$ and $\mu_{j-1} > \mu_j$ if $j > 1$. It follows that $j \leq p-2$, or $j = p-1$, or $j > i$. Since the coordinates are in \mathbb{F}_2 the required verification is trivial. If $\lambda_p = 2$ the proof is similar.

Case (B). $I(\pi) = \{p\} \cup I(\rho)$ and $p \notin I(\rho)$. Then we have $p \notin B(\lambda)$, so $a_p = 0$ and the element $a \in A'(x)$ leaves F_1 and F_{s-1} invariant if and only if $a_p = a_{p+1}$. Assume $a_p = a_{p+1} = 0$. Again using Remark (5.3) one verifies that $a' = a$. If $j \in I(\pi)$ and $j \neq p$ then $j \leq p-2$ or $j \geq p+2$. So the induction is immediate.

7.5. The proof of (7.1) and the verification of (1.2) in our case.

Since H acts transitive on the set of parabolic subgroups of G with a given flag type, cf. Lemma (4.4), Lemma (7.3) implies that every parabolic subgroup of G admits Richardson elements, cf. (1.2)(a). The same argument shows that every polarization of x is obtained by a construction as in (7.3), compare the proof of (1.3). This proves (7.1)(a). Now Lemma (7.4) implies (7.1)(b).

By (4.6)(c), the assertion (7.1)(c) is only then non-trivial if $\lambda = S(\pi)$ and $q = \varepsilon = 0$ and $v^i \equiv 0$ for all i . Then $B(\lambda)$ is empty, cf. (6.6), so that the H -orbit of x splits into two G -orbits, cf. (5.4). Let (p_1, \dots, p_s) be as in (7.3). Put $t = \frac{1}{2}s = v_1$. The construction of (7.3) yields that F_t is generated by the vectors $e(i, j)$ with $i \leq \frac{1}{2}\lambda_j$. So we have

$$F_t = \sum_m x^m \text{Ker}(x^{2m})$$

Since this space is independent of the choice of (p_1, \dots, p_s) the result (7.1)(c) follows easily. Now one may verify (1.2)(b).

Assume that $\lambda = S(\pi)$. By (1.2) the number N_0 of (7.1)(d) can be obtained from (4.6) taking into account the splitting of the Levi type, cf. (4.6) and (6.6)(a). The number N_1 is equal to the index of $A_p(x)$ in $A(x)$. So we have $N_1 = 2^v$ where v is the dimension of the vector space $A(x)/A_p(x)$. If $j \in I(\pi)$ then $j \not\equiv n$ and $j+1 \in B(\lambda)$. This implies that

$$\dim(A'(x)/A'_p(x)) = \# I(\pi) = u$$

in the notations of (7.4). It follows that

$$v \neq u \Leftrightarrow v = u - 1 \Leftrightarrow A'_F(x) \subset A(x) \neq A'(x).$$

We have $A(x) = A'(x)$ if and only if $\varepsilon = 1$ or $B(\lambda)$ is empty. So assume that $\varepsilon = 0$ and $B(\lambda)$ is non-empty. Using (6.6)(b) one verifies that $A'_F(x)$ is not contained in $A(x)$ if and only if $q > 0$.

7.6. Corollary. *For $i = 1, 2$ let $P(i)$ be a polarization of x , say with Levi factor $L(i)$. Let $N_1(i)$ be the number of polarizations of x conjugate to $P(i)$ under $Z_G(x)$. The following conditions are equivalent.*

- (a) $L(1)$ and $L(2)$ are conjugate in G .
- (b) $A_{P(1)}(x) = A_{P(2)}(x)$.
- (c) $N_1(1) = N_1(2)$.

Proof. (a) \Rightarrow (b) by (7.1)(b). The implication (b) \Rightarrow (c) is trivial.

Assume that (c) holds. Write $N_1(1) = N_1(2) = 2^n$. Let $\pi(i) = (v(i); q(i))$ be the Levi type of $P(i)$, so that $\lambda = S(\pi(1)) = S(\pi(2))$. Put $u(i) = \# I(\pi(i))$. By (7.1)(d) we have

$$\begin{aligned} u(i) &= v & \text{if } q(i) + \varepsilon \geq 1 \text{ or } B(\lambda) = \emptyset, \\ u(i) &= v + 1 & \text{if } q(i) = \varepsilon = 0 \text{ and } B(\lambda) \neq \emptyset. \end{aligned}$$

Suppose that $q(1) \neq q(2)$, say $q(1) < q(2)$. By (6.3)(b) we have

$$u(1) - u(2) = \frac{1}{2}(-1)^\varepsilon(q(2) - q(1)) \neq 0.$$

It follows that $\varepsilon = 0$ and $q(1) = 0$ and $u(1) - u(2) = 1$, so that $q(2) = 2$, contradicting the fact that $q(2)$ should be admissible, cf. (6.1). This proves that $q(1) = q(2)$. By (6.5)(a) it follows that $\pi(1) = \pi(2)$ so that (a) follows from (7.1)(c).

7.7. A polarization P of x is called *stable* if P contains $Z_G(x)$. This concept seems to be important in the applications, cf. [2] (6.7)(3).

Corollary. *The element x has stable polarizations if and only if either (i) $\lambda \in \text{Pai}(n)$, or (ii) $n \equiv \varepsilon = 0$ and $\{j \mid \lambda_j \equiv 1\} = \{r, r+1\}$ for some $r \equiv 1$. Assume (i) or (ii) holds. The stable polarizations of x are the N_0 elements of $\text{Pol}(x, \pi)$ where in case (i) we have $\pi = \lambda$, and in case (ii) we have $\pi_j = \lambda_j$ if $j \notin \{r, r+1\}$ and $\pi_r = \lambda_r + 1$ and $\pi_{r+1} = \lambda_{r+1} - 1$.*

The proof is left to the reader. Note that the conditions (i) and (ii) are mutually exclusive.

Remark. If x is an even nilpotent element, cf. [10] page 241, the parabolic subgroup P considered in loc.cit. is a stable polarization of x . The Levi type of P and hence of all stable polarizations of x , can be deduced from the weighted Dynkin diagram of x , cf. loc.cit. page 243, 263 and 264.

7.8. Corollary. *Assume $n \not\equiv \varepsilon$ (i.e. G is of type B_l or C_l).*

- (a) *The polarizations of x have Levi factors with the same semisimple rank.*
- (b) *If P_1 and P_2 are parabolic groups such that their Levi factors have the same Dynkin diagram, then their Richardson elements are conjugate.*

The proof is left to the reader. The table for D_5 in (7.9) clearly shows that both assertions are false for type D_l .

7.9. Tables. A conjugacy class of Levi factors is characterized by the Dynkin type in terms of A_l, B_l, C_l, D_l . The symbols B_1, C_1, D_2, D_3 are used to distinguish certain parts of the Dynkin diagram of G , though formally $B_1 = C_1 = A_1$, $D_2 = A_1 + A_1$, $D_3 = A_3$. The same information is carried by the Levi type $(v; q)$. The partition v is given by the sequence $v^* = (v^1, \dots, v^t)$ where $v^t > 0 = v^{t+1}$.

The numbers N_0 and N_1 are meant conform (1.4) and (7.1). The partition $\lambda = S(\pi)$ of a Richardson element x is given by the sequence $\lambda_* = (\lambda_1, \dots, \lambda_m)$ where $\lambda_m > 0 = \lambda_{m+1}$. The groups $A(x)$ and $A_p(x)$ are determined by the sets $B(\lambda)$ and $I(\pi)$, cf. (5.3) and (7.1). These sets are given in the tables.

We give the weighted Dynkin diagram of x , in view of remark (7.7). If the H -orbit of x or equivalently the Levi type, splits into two conjugacy classes under the action of G , we write $2 \times N_0$ in the column of N_0 (so that this column adds up to 2^l). In this case we give the Dynkin diagram of one of the G -orbits. We indicate how to get the other diagram by the symbol \curvearrowright . A partition λ such that the corresponding nilpotent elements have no polarizations at all, is equipped with a dot and dash line in the column of (v^*, q) .

Tabelle 1




$B_2 = \text{SO}(5)$							
Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	(1 1; 1)	1	1	\emptyset	1	5	2 2
B_1	(1; 3)	1	1	\emptyset	1 3	3 1 1	2 0
A_1	(2; 1)	1	2	2			
	— · — · —				3	2 2 1	0 1
B_2	(—; 5)	1	1	\emptyset	5	1^5	0 0
$C_2 = \text{Sp}(4)$							
Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	(1 1; 0)	1	1	\emptyset	1	4	2 2
A_1	(2; 0)	1	1	\emptyset	2	2 2	0 2
C_1	(1; 2)	1	2	1			
	— · — · —				1	2 1 1	1 0
C_2	(—; 4)	1	1	\emptyset	\emptyset	1^4	0 0
$B_3 = \text{SO}(7)$							
Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	(1 1 1; 1)	1	1	\emptyset	1	7	2 2 2
B_1	(1 1; 3)	1	1	\emptyset	1 3	5 1 1	2 2 0
A_1	(2 1; 1)	2	2	2			
$A_1 + B_1$	(2; 3)	1	1	\emptyset	2 3	3 3 1	0 2 0
A_2	(3; 1)	1	1	\emptyset	1	3 2 2	1 0 1
B_2	(1; 5)	1	1	\emptyset	1 5	3 1^4	2 0 0
	— · — · —				5	2 2 1^3	0 1 0
B_3	(—; 7)	1	1	\emptyset	7	1^7	0 0 0

Table 1 (continued)

 $C_3 = \text{Sp}(6)$

Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	$\bigcirc - \bigcirc - \bigcirc - \bigcirc$
0	(1 1 1; 0)	1	1	\emptyset	1	6	2 2 2
A_1	(2 1; 0)	2	1	$\emptyset \}$	1 2	4 2	2 0 2
C_1	(1 1; 2)	1	2	1 }			
	--- --				1	4 1 1	2 1 0
$A_1 + C_1$	(2; 2)	1	1	\emptyset	\emptyset	3 3	0 2 0
A_2	(3; 0)	1	1	\emptyset	3	2 2 2	0 0 2
C_2	(1; 4)	1	2	1	2	2 2 1 1	0 1 0
	--- --				1	2 1 ⁴	1 0 0
C_3	(-; 6)	1	1	\emptyset	6	1 ⁶	0 0 0

 $B_4 = \text{SO}(9)$


Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	$\bigcirc - \bigcirc - \bigcirc - \bigcirc - \bigcirc$
0	(1 1 1 1; 1)	1	1	\emptyset	1	9	2 2 2 2
B_1	(1 1 1; 3)	1	1	$\emptyset \}$	1 3	7 1 1	2 2 2 0
A_1	(2 1 1; 1)	3	2	2 }			
$A_1 + B_1$	(2 1; 3)	2	1	$\emptyset \}$	1 2 3	5 3 1	2 0 2 0
$A_1 + A_1$	(2 2; 1)	1	2	2 }			
A_2	(3 1; 1)	2	1	\emptyset	1	5 2 2	2 1 0 1
B_2	(1 1; 5)	1	1	\emptyset	1 5	5 1 ⁴	2 2 0 0
	--- --				3	4 4 1	0 2 0 1
$A_2 + B_1$	(3; 3)	1	1	\emptyset	3	3 3 3	0 0 2 0
$A_1 + B_2$	(2; 5)	1	1	\emptyset	2 5	3 3 1 ³	0 2 0 0
A_3	(4; 1)	1	2	4	1 5	3 2 2 1 1	1 0 1 0
B_3	(1; 7)	1	1	\emptyset	1 7	3 1 ⁶	2 0 0 0
	--- --				5	2 ⁴ 1	0 0 0 1
	--- --				7	2 2 1 ⁵	0 1 0 0
B_4	(-; 9)	1	1	\emptyset	9	1 ⁹	0 0 0 0

 $C_4 = \text{Sp}(8)$


Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	$\bigcirc - \bigcirc - \bigcirc - \bigcirc - \bigcirc$
0	(1 1 1 1; 0)	1	1	\emptyset	1	8	2 2 2 2
A_1	(2 1 1; 0)	3	1	$\emptyset \}$	1 2	6 2	2 2 0 2
C_1	(1 1 1; 2)	1	2	1 }			
	--- --				1	6 1 1	2 2 1 0
$A_1 + A_1$	(2 2; 0)	1	1	$\emptyset \}$	2	4 4	0 2 0 2
$A_1 + C_1$	(2 1; 2)	2	2	1 }			
A_2	(3 1; 0)	2	1	\emptyset	1 3	4 2 2	2 0 0 2
C_2	(1 1; 4)	1	2	1	1 2	4 2 1 1	2 0 1 0
	--- --				1	4 1 ⁴	2 1 0 0
$A_2 + C_1$	(3; 2)	1	1	\emptyset	3	3 3 2	0 1 1 0
$A_1 + C_2$	(2; 4)	1	1	\emptyset	\emptyset	3 3 1 1	0 2 0 0
A_3	(4; 0)	1	1	\emptyset	4	2 ⁴	0 0 0 2
	--- --				3	2 ³ 1 1	0 0 1 0
C_3	(1; 6)	1	2	1	2	2 2 1 ⁴	0 1 0 0
	--- --				1	2 1 ⁶	1 0 0 0
C_4	(-; 8)	1	1	\emptyset	\emptyset	1 ⁸	0 0 0 0

Table 1 (continued)

 $D_4 = \text{SO}(8)$

Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	$(1 \ 1 \ 1 \ 1; 0)$	1	1	1	1 2	7 1	2 2 2 2
A_1	$(2 \ 1 \ 1; 0)$	4	1	1	1 2	5 3	2 0 2 2
$A_1 + A_1$	$(2 \ 2; 0)$	2×1	1	\emptyset	\emptyset	4 4	0 2 $2 \rightsquigarrow 0$
D_2	$(1 \ 1; 4)$	1	1	\emptyset	1 4	5 1 1 1	2 2 0 0
$A_1 + D_2$	$(2; 4)$	1	1	\emptyset	2 4	3 3 1 1	0 2 0 0
A_2	$(3 \ 1; 0)$	3	2	1 3			
---	---	---	---	---	1 4	3 2 2 1	1 0 1 1
A_3	$(4; 0)$	2×1	1	\emptyset	\emptyset	2 2 2 2	0 0 $2 \rightsquigarrow 0$
D_3	$(1; 6)$	1	1	\emptyset	1 6	3 1 ⁵	2 0 0 0
---	---	---	---	---	6	2 2 1 ⁴	0 1 0 0
D_4	$(-; 8)$	1	1	\emptyset	8	1 ⁸	0 0 0 0

 $D_5 = \text{SO}(10)$

Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	$(1^5; 0)$	1	1	1	1 2	9 1	2 2 2 2 2
A_1	$(2 \ 1 \ 1 \ 1; 0)$	5	1	1	1 2	7 3	2 2 0 2 2
D_2	$(1 \ 1 \ 1; 4)$	1	1	\emptyset	1 4	7 1 1 1	2 2 2 0 0
$A_1 + A_1$	$(2 \ 2 \ 1; 0)$	5	1	1	2	5 5	0 2 0 2 2
$A_1 + D_2$	$(2 \ 1; 4)$	2	1	\emptyset	1 2 4	5 3 1 1	2 0 2 0 0
A_2	$(3 \ 1 \ 1; 0)$	4	2	1 3			
---	---	---	---	---	1 4	5 2 2 1	2 1 0 1 1
D_3	$(1 \ 1; 6)$	1	1	\emptyset	1 6	5 1 ⁵	2 2 0 0 0
$A_1 + A_2$	$(3 \ 2; 0)$	4	1	3	4	4 4 1 1	0 2 0 1 1
$A_2 + D_2$	$(3; 4)$	1	1	\emptyset	3 4	3 3 3 1	0 0 2 0 0
A_3	$(4 \ 1; 0)$	3	1	1	2	3 3 2 2	0 1 0 1 1
$A_1 + D_3$	$(2; 6)$	1	1	\emptyset	2 6	3 3 1 ⁴	0 2 0 0 0
---	---	---	---	---	1 6	3 2 2 1 ³	1 0 1 0 0
D_4	$(1; 8)$	1	1	\emptyset	1 8	3 1 ⁷	2 0 0 0 0
A_4	$(5; 0)$	2	1	5	6	2 ⁴ 1 1	0 0 0 1 1
---	---	---	---	---	8	2 2 1 ⁶	0 1 0 0 0
D_5	$(-; 10)$	1	1	\emptyset	10	1 ¹⁰	0 0 0 0 0

 $D_6 = \text{SO}(12)$


Levi	$(v^*; q)$	N_0	N_1	$I(\pi)$	$B(\lambda)$	λ_*	
0	$(1^6; 0)$	1	1	1	1 2	11, 1	2 2 2 2 2 2
A_1	$(2 \ 1^4; 0)$	6	1	1	1 2	9 3	2 2 2 0 2 2
D_2	$(1^4; 4)$	1	1	\emptyset	1 4	9 1 1 1	2 2 2 2 0 0
$A_1 + A_1$	$(2 \ 2 \ 1 \ 1; 0)$	9	1	1	1 2	7 5	2 0 2 0 2 2
$A_1 + D_2$	$(2 \ 1 \ 1; 4)$	3	1	\emptyset	1 2 4	7 3 1 1	2 2 0 2 0 0
A_2	$(3 \ 1 \ 1 \ 1; 0)$	5	2	1 3			
---	---	---	---	---	1 4	7 2 2 1	2 2 1 0 1 1
D_3	$(1 \ 1 \ 1; 6)$	1	1	\emptyset	1 6	7 1 ⁵	2 2 2 0 0 0
$3A_1$	$(2 \ 2 \ 2; 0)$	2×1	1	\emptyset	\emptyset	6 6	0 2 0 2 $2 \rightsquigarrow 0$
$2A_1 + D_2$	$(2 \ 2; 4)$	2	1	\emptyset	2 4	5 5 1 1	0 2 0 2 0 0
$A_1 + A_2$	$(3 \ 2 \ 1; 0)$	10	2	1 3			

Table 1 (continued)

$A_2 + D_2$	(3 1; 4)	2	1	\emptyset	1 3 4	5 3 3 1	2 0 0 2 0 0
A_3	(4 1 1; 0)	4	1	1	1 2	5 3 2 2	2 0 1 0 1 1
$A_1 + D_3$	(2 1; 6)	2	1	\emptyset	1 2 6	5 3 1 ⁴	2 0 2 0 0 0
	— — — —				1 6	5 2 2 1 ³	2 1 0 1 0 0
D_4	(1 1; 8)	1	1	\emptyset	1 8	5 1 ⁷	2 2 0 0 0 0
$A_2 + A_2$	(3 3; 0)	2	1	3	3 4	4 4 3 1	0 1 1 0 1 1
$A_3 + A_1$	(4 2; 0)	2×2	1	\emptyset	\emptyset	4 4 2 2	0 2 0 0 2 \curvearrowright 0
	— — — —				6	4 4 1 ⁴	0 2 0 1 0 0
$A_3 + D_2$	(4; 4)	1	1	\emptyset	4	3 3 3 3	0 0 0 2 0 0
$A_2 + D_3$	(3; 6)	1	1	\emptyset	3 6	3 ³ 1 ³	0 0 2 0 0 0
A_4	(5 1; 0)	3	2	1 5	2 6	3 3 2 2 1 1	0 1 0 1 0 0
$A_1 + D_4$	(2; 8)	1	1	\emptyset	2 8	3 3 1 ⁶	0 2 0 0 0 0
	— — — —				1 6	3 2 2 2 2 1	1 0 0 0 1 1
	— — — —				1 8	3 2 2 1 ⁵	1 0 1 0 0 0
D_5	(1; 10)	1	1	\emptyset	1, 10	3 1 ⁹	2 0 0 0 0 0
A_5	(6; 0)	2×1	1	\emptyset	\emptyset	2 ⁶	0 0 0 0 2 \curvearrowright 0
	— — — —				8	2 ⁴ 1 ⁴	0 0 0 1 0 0
	— — — —				10	2 2 1 ⁸	0 1 0 0 0 0
D_6	(—; 12)	1	1	\emptyset	12	1 ¹²	0 0 0 0 0 0

References

1. Borho, W.: Recent advances in enveloping algebras of semi-simple Lie-algebras. Séminaire Bourbaki, 29e année, 1976/77, n° 489. Berlin-Heidelberg-New York: Springer (to appear)
2. Borho, W.: Definition einer Dixmier-Abbildung für $\mathfrak{sl}(n, \mathbb{C})$. *Inventiones math.* **40**, 143–169 (1977)
3. Borho, W., Jantzen, J.C.: Über primitive Ideale in der Einhüllenden einer halbeinfachen Lie-algebra. *Inventiones math.* **39**, 1–53 (1977)
4. Borho, W., Kraft, H.: Über Bahnen und deren Deformationen bei linearen Aktionen reduktiver Gruppen. Preprint
5. Hesselink, W.H.: Singularities in the nilpotent scheme of a classical group. *Trans. Amer. Math. Soc.* **222**, 1–32 (1976)
6. Humphreys, J.E.: Linear algebraic groups. Berlin-Heidelberg-New York: Springer 1975
7. Ozeki, H., Wakimoto, M.: On polarizations of certain homogeneous spaces. *Hiroshima Math. J.* **2**, 445–482 (1972)
8. Richardson, R.W.: Conjugacy classes in parabolic subgroups of semi-simple algebraic groups. *Bull. London Math. Soc.* **6**, 21–24 (1974)
9. Spaltenstein, N.: Sous-groupes de Borel contenant un unipotent donné. Preprint
10. Springer, T.A., Steinberg, R.: Conjugacy classes. In: Borel, A. et al.: Seminar on algebraic groups and related finite groups. *Lecture Notes in Mathematics* 131, pp. 167–266. Berlin-Heidelberg-New York: Springer 1970
11. Steinberg, R.: Conjugacy classes in algebraic groups. *Lecture Notes in Mathematics* 366. Berlin-Heidelberg-New York: Springer 1974

Received November 21, 1977